



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Now, the expression e^{2ax^2} can be integrated only, so far as we know, by expanding it in an infinite series, thus:

$$e^{2ax^2} = 1 + 2ax^2 + \frac{4a^2x^4}{1.2} + \dots$$

$$\therefore \int e^{2ax^2} dx = x + \frac{2ax^3}{1.3} + \frac{4a^2x^5}{1.2.5} + \dots$$

NOTE. Professors Harding and Prime should have received credit for solving 308.

PROBLEMS FOR SOLUTION.

ALGEBRA.

366. Proposed by WILLIAM HOOVER, Ph. D., Professor of Mathematics and Astronomy, Ohio University, Athens, Ohio.

Eliminate m from the equations

$$\begin{aligned} 3a^2m^4 + 4aym^3 + 6axm^2 - (x^2 + y^2 - 4ax) &= 0; \\ (x^2 + y^2 - 4ax)m^4 - 6axm^2 + 4aym - 3a^2 &= 0. \end{aligned}$$

367. Proposed by E. B. ESCOTT, University of Michigan, Ann Arbor, Michigan.

Solve the simultaneous equations:

$$\frac{2x}{1+x^2} = y \dots (1); \quad \frac{2y}{1+y^2} = z \dots (2); \quad \frac{2z}{1+z^2} = u \dots (3); \quad \frac{2u}{1+u^2} = x \dots (4).$$

368. Proposed by E. B. ESCOTT, University of Michigan, Ann Arbor, Michigan.

Solve the functional equation, $\frac{f(-x)}{f(x)} = r^{2x}$.

GEOMETRY.

399. Proposed by J. K. ELLWOOD, Superintendent of Schools, Lucas, Kansas.

A race track is to be composed of two tangents and the arc of the circle which is concave towards the point of intersection of the two tangents, each tangent and the arc of the circle being 1 mile. What is the radius of the circle?

400. Proposed by FRANCIS RUST, C. E., Pittsburgh, Pennsylvania.

Given a circle and a point P without; construct, *using the straight edge only*, the two tangents to the circle through P .